

# Educational resources on cervical dilation with the Cervical Ripening Balloon

1. Alfirevic Z, Keeney E, Dowswell T, et al. Methods to induce labour: a systematic review, network meta-analysis and cost-effectiveness analysis. *BJOG*. 2016;123(9):1462-1470.
2. Cromi A, Ghezzi F, Uccella S, et al. A randomized trial of preinduction cervical ripening: dinoprostone vaginal insert versus double-balloon catheter. *Am J Obstet Gynecol*. 2012;207(2):125.e1-e7. doi:10.1016/j.ajog.2012.05.020.
3. Du C, Liu Y, Liu Y, et al. Double-balloon catheter vs. dinoprostone vaginal insert for induction of labor with an unfavorable cervix. *Arch Gynecol Obstet*. 2015;291(6):1221-1227.
4. Du YM, Zhu LY, Cui LN, et al. Double-balloon catheter versus prostaglandin E2 for cervical ripening and labour induction: a systematic review and meta-analysis of randomised controlled trials [published online ahead of print August 17, 2016]. *BJOG*. 2016. doi: 10.1111/1471-0528.14256.
5. Hoppe KK, Schiff MA, Peterson SE, et al. 30 mL single- versus 80 mL double-balloon catheter for pre-induction cervical ripening: a randomized controlled trial [published online ahead of print August 25, 2015]. *J Matern Fetal Neonatal Med*. 2015. doi:10.3109/14767058.2015.1067297.
6. Mei-Dan E, Walfisch A, Suarez-Easton S, et al. Comparison of two mechanical devices for cervical ripening: a prospective quasi-randomized trial. *J Matern Fetal Neonatal Med*. 2012;25(6):723-727.
7. National Institute for Health and Care Excellence. Insertion of a double balloon catheter for induction of labour in pregnant women without previous caesarean section. NICE Web site. <https://www.nice.org.uk/guidance/ipg528>. Published July 2015. Accessed October 26, 2016.
8. Pennell CE, Henderson JJ, O'Neill MJ, et al. Induction of labour in nulliparous women with an unfavourable cervix: a randomised controlled trial comparing double and single balloon catheters and PGE2 gel. *BJOG*. 2009;116(11):1443-1452.
9. Salim R, Zafran N, Nachum Z, et al. Single-balloon compared with double-balloon catheters for induction of labor: a randomized controlled trial. *Obstet Gynecol*. 2011;118(1):79-86.
10. Shechter-Maor G, Haran G, Sadeh-Mestechkin D, et al. Intra-vaginal prostaglandin E2 versus double-balloon catheter for labor induction in term oligohydramnios. *J Perinatol*. 2015;35(2):95-98.
11. Suffecool K, BM Rosenn, Kam S, et al. Labor induction in nulliparous women with an unfavorable cervix: double balloon catheter versus dinoprostone. *J Perinat Med*. 2014;42(2):213-218.
12. Tan TL, Ng GYH, Lim SEL, et al. Cervical ripening balloon as an alternative for induction of labour: a randomized controlled trial. *BJMP*. 2015;8(1):a806.